arXiv:math/0507193v1 [math.PR] 10 Jul 2005

Lévy processes:
Hitting Time, Overshoot and Undershoot
II - Asymptotic Behaviour

Bernard ROYNETTE$^{(1)}$, **Pierre VALLOIS**$^{(1)}$ and **Agnès VOLPI** $^{(2)}$

Manuscript P661 submitted to SPA, October 2004

(1) Université Henri Poincaré, Institut de Mathématiques Elie Cartan, B.P. 239, F-54506 Vandœuvre-lès-Nancy Cedex
(2) ESSTIN, 2 rue Jean Lamour, Parc Robert Bentz, 54500 Vandœuvre-lès-Nancy, France.
e-mail:
roynette@iecn.u-nancy.fr vallois@iecn.u-nancy.fr agnes.volpi@esstin.uhp-nancy.fr

Abstract

Let $(X_t,\ t \geq 0)$ be a Lévy process started at 0, with Lévy measure ν and T_x the first hitting time of level $x > 0$: $T_x := \inf\{t \geq 0;\ X_t > x\}$. Let $F(\theta, \mu, \rho, .)$ be the joint Laplace transform of (T_x, K_x, L_x): $F(\theta, \mu, \rho, x) := \mathbb{E}\left(e^{-\theta T_x - \mu K_x - \rho L_x} \mathbb{1}_{\{T_x < +\infty\}}\right)$, where $\theta \geq 0$, $\mu \geq 0$, $\rho \geq 0$, $x \geq 0$, $K_x := X_{T_x} - x$ and $L_x := x - X_{T_x -}$.
If we assume that ν has finite exponential moments we exhibit an asymptotic expansion for $F(\theta, \mu, \rho, x)$, as $x \to +\infty$. A limit theorem involving a normalization of the triplet (T_x, K_x, L_x) as $x \to +\infty$, may be deduced.
At last, if $\nu_{|\mathbb{R}_+}$ has finite moment of fixed order, we prove that the ruin probability $\mathbb{P}(T_x < +\infty)$ has at most a polynomial decay.

Keywords: Lévy processes, ruin problem, hitting time, overshoot, undershoot, asymptotic estimates, functional equation, characteristic function, exponential and polynomial decay.

AMS 2000 Subject classification: 60E10, 60F05, 60G17, 60G40, 60G51,60J35, 60J65, 60J75, 60J80.

1 Introduction

1.1 Let $(X_t\ ,\ t \geq 0)$ be a Lévy process started at 0, with decomposition :

$$X_t = \sigma B_t - c_0 t + J_t \quad ; \quad t \geq 0,$$

where $c_0 \in \mathbb{R}$, $\sigma > 0$, $(B_t\,,\ t\geq 0)$ is a standard Brownian motion started at 0, $(J_t\,,\ t\geq 0)$ is a pure jump process, independent of $(B_t\,,\ t\geq 0)$.
Let us denote by ν its Lévy measure, and T_x the first hitting time of level $x > 0$:

$$T_x := \inf\{t \geq 0\ ;\ X_t > x\}\,. \tag{1.1}$$

In this paper we essentially focus on the asymptotic behaviour of T_x as $x \to \infty$. Since $T_x = \inf\{t \geq 0; X_t/\sigma > x/\sigma\}$, we can assume that $\sigma = 1$. Consequently, we have :

$$X_t = B_t - c_0 t + J_t \quad ; \quad t \geq 0. \tag{1.2}$$

If $(X_t\,,\ t\geq 0)$ is a Brownian motion with drift $-c_0$ (i.e. $J_t = 0$), then (cf. [8] p.197) :

$$P(T_x < +\infty) = \begin{cases} 1 & \text{if } c_0 \leq 0 \\ e^{-2c_0 x} & \text{if } c_0 > 0. \end{cases} \tag{1.3}$$

In particular, if $c_0 > 0$, there exists $\gamma > 0$, $C, C' > 0$ such that

$$\mathbb{P}(T_x < +\infty) \leq C' e^{-\gamma x} \tag{1.4}$$

$$\mathbb{P}(T_x < +\infty) \underset{x\to+\infty}{\sim} C e^{-\gamma x}\,. \tag{1.5}$$

A lot of authors have generalized (1.4) when $(X_t\,,\ t\geq 0)$ is either a Lévy process, or a diffusion, see for instance [6], [5], [13], [14], [3], [9].
Suppose that $(X_t\,,\ t\geq 0)$ is a Lévy process with no negative jump (i.e. the support of its Lévy measure is included in $\mathbb{R}_+$) and the characteristic exponent ψ of $(X_t\,,\ t\geq 0))$ has a positive zero γ_0. Then, in [2], it has proved that (1.5) holds, with $\gamma = \gamma_0$ and $C = -\dfrac{\psi'(0)}{\psi'(\gamma_0)}$. This result was extended in [1], without any restriction on the support of ν. However in this case C is not explicit.

1.2 Since (X_t) may jump, if $T_x < \infty$, we have : $X_{T_x} \geq x \geq X_{T_x-}$, where X_{T_x-} denotes the left-limit of (X_t) at time T_x. Therefore it is interesting to consider the overshoot K_x and the undershoot L_x, these r.v.'s being defined on $\{T_x < +\infty\}$ as follows :

$$K_x := X_{T_x} - x\ ; \qquad L_x := x - X_{T_x-}\,. \tag{1.6}$$

One aim of this paper is a generalization of (1.5). We first replace $\mathbb{P}(T_x<+\infty)$ by $F(\theta,\mu,\rho,x)$, where $F(\theta,\mu,\rho,\cdot)$ is the Laplace transform of the triplet (T_x, K_x, L_x) :

$$F(\theta,\mu,\rho,x) := \mathbb{E}\left(e^{-\theta T_x - \mu K_x - \rho L_x} \mathbb{1}_{\{T_x<+\infty\}}\right)\,. \tag{1.7}$$

Second, we do not restrict ourselves to determine an equivalent of $F(\theta,\mu,\rho,x)$, as x goes to infinity, we obtain (cf Theorem 2.4) an asymptotic development of this function. In our proof, the x-Laplace transform :

$$\widehat{F}(\theta,\mu,\rho,q) := \int_0^{+\infty} e^{-qx} F(\theta,\mu,\rho,x)dx\,, \tag{1.8}$$

of $F(\theta,\mu,\rho,.)$ plays a central role.
Let us explain the idea of our approach. For simplicity we first consider $\mathbb{P}(T_x<+\infty)$ (i.e. $\theta=\mu=\rho=0$).
Obviously (1.5) is equivalent to $\lim\limits_{x\to+\infty} e^{\gamma x}\mathbb{P}(T_x<+\infty)=C$. Suppose that condition holds, then

$$\int_0^{+\infty} e^{-qx}e^{\gamma x}\mathbb{P}(T_x<+\infty)dx=\widehat{F}(0,0,0,q-\gamma)\underset{q\to 0}{\sim}\frac{C}{q}\,. \tag{1.9}$$

Hence C is the residue of $\widehat{F}(0,0,0,q-\gamma)$ at 0.
To deal with $F(\theta,\mu,\rho,x)$, we have to suppose that the jumps of (X_t) are not too big :

$$\int_{-\infty}^{-1} e^{-qy}\nu(dy)<+\infty \quad \forall q>0\,, \tag{1.10}$$

and

$$\int_1^{\infty} e^{sy}\nu(dy)<+\infty \quad \text{for some } \; s>0\,. \tag{1.11}$$

With some additional assumptions (i.e. (2.7) and (2.16)), we prove in Theorem 2.4 :

$$\begin{aligned} F(\theta,\mu,\rho,x)=&C_0(\theta,\mu,\rho)e^{-\gamma_0(\theta)x}+\sum_{i=1}^{p} a_i\left(C_i(\theta,\mu,\rho,x)e^{-\gamma_i(\theta)x}+\overline{C_i}(\theta,\mu,\rho,x)e^{-\overline{\gamma_i}(\theta)x}\right)\\ &+\mathrm{O}\left(e^{-Bx}\right)\,, \end{aligned} \tag{1.12}$$

where $C_0(\theta,\mu,\rho)$ is a positive real number, $C_1(\theta,\mu,\rho,x),\cdots,C_p(\theta,\mu,\rho,x)$ are x-polynomial functions with values in $\mathbb{C}$, $(\gamma_0(\theta),\gamma_1(\theta),\cdots,\gamma_p(\theta),\overline{\gamma_1}(\theta),\cdots,\overline{\gamma_p}(\theta))$ are zeros of $\varphi-\theta$ (where φ is the function defined by (2.1)) and $a_i=\dfrac{1}{2}$ (resp. 1) if $\gamma_i(\theta)\in\mathbb{R}$ (otherwise).
Few words about the proof of (1.12). Starting with a functional equation verified by $\widehat{F}(\theta,\mu,\rho,.)$ ([12],Theorem 2.1), we determine the poles and the residues of $q\mapsto\widehat{F}(\theta,\mu,\rho,q-\gamma_0(\theta))$. Then, using the Mellin-Fourier inverse transformation we can recover $F(\theta,\mu,\rho,.)$.
1.3 In the same spirit, we investigate polynomial decay of the probability of ruin. More precisely we prove in Theorem 3.1 :

$$P(T_x<\infty)\le\frac{C}{1+x^n},\quad \forall x\ge 0, \tag{1.13}$$

where it is supposed that $\int_0^{\infty} y^p\nu(dy)<\infty$, for some $p\ge 2$, n being the integer part of $p-2$.
Some results of this type have been obtained in [11], [4], when $(X_t\,,\;t\ge 0)$ is a diffusion .

1.4 In section 4 we give two applications of Theorem 2.4. The first one concerns the asymptotic behaviour of the triplet (T_x, K_x, L_x) as $x \to \infty$. Since $x \to T_x$ is non-decreasing, the first component T_x has to be normalized in $\widehat{T}_x$. We prove in Theorem 4.1 (see also Remark 4.2) that $(\widehat{T}_x, K_x, L_x)$ converges in distribution, as $x \to +\infty$. The first component $\widehat{T}_x$ converges to a Gaussian distribution. Morover time and position become asymptotically independent. In [9], the limit of distribution of the overshoot has been investigated. A. Gut [7] has also studied the convergence of the triplet, the Lévy process $(X_t\,,\ t \geq 0)$ being replaced by a random walk.

The asymptotic development of $F(\theta, \mu, \rho, x)$ given in Theorem 2.4 gives the rate of convergence of $\widehat{T}_x$ to the Gaussian distribution (see Remark 4.3 for details).

2 Asymptotic expansion of $F(\theta, \mu, \rho, .)$

We keep the notation given in the Introduction. Let ψ the characteristic exponent of $(X_t\,,\ t \geq 0)$, i.e. $\mathbb{E}\left(e^{qX_t}\right) = e^{t\psi(q)}$. For our purpose it is more convenient to deal with the function φ, where $\varphi(q) = \psi(-q)$. Lévy Khintchine formula gives :

$$\varphi(q) = \frac{q^2}{2} + cq + \int_{\mathbb{R}} \left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}}\right) \nu(dy)\,. \tag{2.1}$$

In this section, it is supposed that ν verifies :

$$r_\nu > 0 \quad \text{(may be} \quad r_\nu = +\infty)\,, \tag{2.2}$$

$$r_\nu^* = \infty\,, \tag{2.3}$$

where

$$r_\nu := \sup\left\{s \geq 0;\ \int_1^{+\infty} e^{sy}\nu(dy)\ <\ +\infty\right\} \in [0, +\infty]\,, \tag{2.4}$$

$$r_\nu^* := \sup\ \left\{q \in \mathbb{R}\,;\ \int_{-\infty}^{-1} e^{-qy}\nu(dy) < +\infty\right\}, \tag{2.5}$$

with the convention $\sup \emptyset = 0$.

Note that (2.3) is equivalent to (1.10).

Consequently, in our setting, φ is defined on $]-r_\nu, +\infty[$ and $\displaystyle\int_{|y|>1} |y|\nu(dy) < \infty$.

In particular :

$$E[|X_1|] < \infty. \tag{2.6}$$

Assumptions (2.3) and (2.2) will be needed throughout this section, therefore we do not repeat them in the statement of the results.

Let us briefly sketch the proof of our main result (Theorem 2.4). We start with the functional equation verified by $\widehat{F}(\theta,\mu,\rho,.)$. This result was established in [12], and the formula is recalled in (2.33). To recover $F(\theta,\mu,\rho,.)$ we use the Mellin-Fourier inverse transformation. This leads us prove that some modification (namely the function $\widehat{\widehat{F}}(\theta,\mu,\rho,\cdot)$ defined by (2.36)) of $t\to\widehat{F}(\theta,\mu,\rho,q+it)$ is an integrable function. We have also to finely analyze the zeros of $\varphi-\theta$.
As it is briefly explained in 1.2 of the Introduction, the zeros of the function $\varphi_\theta=\varphi-\theta$ play a important role.
Let us start with the negative zeros of φ_θ. We assume (cf. Annex of [12], Figures 3.a, 4.a and 5) that there exists $\kappa>0$, such that for any $\theta\in[0,\kappa]$:

$$\exists\ -\gamma_0(\theta)\in]-r_\nu,0] \quad \text{satisfying} \quad \varphi(-\gamma_0(\theta))=\theta\,. \tag{2.7}$$

More precisely :

$$(i)\quad -\gamma_0(\theta)<0, \quad \text{if} \quad \theta>0\,, \tag{2.8}$$

$$(ii)\quad -\gamma_0(0)<0, \quad \text{if} \quad \theta=0 \quad \text{and} \quad \mathbb{E}(X_1)<0\,, \tag{2.9}$$

$$(iii)\quad -\gamma_0(0)=0, \quad \text{if} \quad \theta=0 \quad \text{and} \quad \mathbb{E}(X_1)\geq 0\,. \tag{2.10}$$

As for the positive zeros of φ_θ, it is proved in Annex of [12] :

$$\forall\,\theta\geq 0, \qquad \exists\,\gamma_0^*(\theta)\geq 0 \qquad \text{such that} \qquad \varphi(\gamma_0^*(\theta))=\theta\,. \tag{2.11}$$

More precisely :

$$(i)\quad \gamma_0^*(\theta)>0, \quad \text{if} \quad \theta>0\,, \tag{2.12}$$

$$(ii)\quad \gamma_0^*(0)>0, \quad \text{if} \quad \theta=0 \quad \text{and} \quad \mathbb{E}(X_1)>0\,, \tag{2.13}$$

$$(iii)\quad \gamma_0^*(0)=0, \quad \text{if} \quad \theta=0 \quad \text{and} \quad \mathbb{E}(X_1)\leq 0\,. \tag{2.14}$$

We set $D_b:=\{q\in\mathbb{C};\quad \text{such that} \quad \operatorname{Re}q>-b\}$ and we introduce :

$$B_\nu:=\sup\{b>0\,;\ \widehat{\nu_{|[1,+\infty[}} \text{ admits a meromorphic extension to } D_b\}\,, \tag{2.15}$$

where $\quad \widehat{\nu_{|[1,+\infty[}}(q):=\displaystyle\int_1^{+\infty}e^{-qy}\nu(dy).$

Thanks to (2.2), we note that $B_\nu\geq r_\nu$ and B_ν may be equal to ∞. Then φ has a meromorphic extension to D_{B_ν}. For simplicity we still denote φ this extension. Likewise, $\widehat{\nu_{|[1,+\infty[}}$ is the meromorphic extension of $\widehat{\nu_{|[1,+\infty[}}$ to D_{B_ν}.
Our last assumption on ν is :

$$\begin{aligned}&\forall B\in]0,B_\nu[,\quad \exists K>0,\quad \exists R_0>0, \text{such that :}\\ &|\widehat{\nu_{|[1,+\infty[}}(q)|\leq K|q|\,,\quad \text{for any } q \text{ such that} \quad \begin{cases}-B\leq \operatorname{Re}q\leq 0\\ |\operatorname{Im}q|\geq R_0\end{cases}\end{aligned} \tag{2.16}$$

A large class of measures ν satisfying previous hypothesis and (2.16) will be given in Remark 2.5 below.

Remark 2.1 *1. Relations (2.2) and (2.3) allow us to replace* $\widehat{\nu_{|[1,+\infty[}}(q)$ *in (2.16), by either* $\int_0^{+\infty}(e^{-qy}-1+qy\mathbb{1}_{\{|y|<1\}})\nu(dy)$ *or* $\int_{-\infty}^{+\infty}(e^{-qy}-1+qy\mathbb{1}_{\{|y|<1\}})\nu(dy)$.

2. Since :

$$\sup_{\mathrm{Re}\, q\geq 0}|\widehat{\nu_{|[1,+\infty[}}(q)|<+\infty\,, \tag{2.17}$$

then (2.16) implies :

$$\forall q \quad \text{such that} \ : \ \begin{cases}\mathrm{Re}\, q\geq -B\\ |\mathrm{Im}\, q|\geq R_0\end{cases} \qquad |\widehat{\nu_{|[1,+\infty[}}(q)|\leq K|q|. \tag{2.18}$$

The key result concerning the complex zeros of φ_θ is the following.

Proposition 2.2 *We suppose that (2.7) and (2.16) hold. Then for any* $\theta\in[0,\kappa]$, *there exists* $\beta_\theta>0$ *such that for any* $B\in]0,B_\nu[$, φ_θ *admits a finite number of conjugated zeros in the strip* $D_{B,\beta_\theta}:=\{q\in\mathbb{C}\ ;\ -B\leq\mathrm{Re}\, q\leq\beta_\theta\}$.
This set of zeros of φ_θ *in* D_{B,β_θ} *is equal to :*

1. $\{-\gamma_0(\theta),\ -\gamma_1(\theta),\ -\overline{\gamma_1}(\theta),\cdots,-\gamma_p(\theta),\ -\overline{\gamma_p}(\theta)\}$ *if* $\theta>0$,
2. $\{0,\ -\gamma_0(0),\ -\gamma_1(0),\ -\overline{\gamma_1}(0),\cdots,-\gamma_p(0),\ -\overline{\gamma_p}(0)\}$ *if* $\theta=0$ *and* $\mathbb{E}(X_1)<0$,
3. $\{-\gamma_0(0)=0,\ -\gamma_1(0),\ -\overline{\gamma_1}(0),\cdots,-\gamma_p(0),\ -\overline{\gamma_p}(0)\}$ *if* $\theta=0$ *and* $\mathbb{E}(X_1)\geq 0$,

   where

$$-B<-\mathrm{Re}\,(\gamma_p(\theta))\leq\cdots\leq-\mathrm{Re}\,(\gamma_1(\theta))<-\gamma_0(\theta)\leq 0\,. \tag{2.19}$$

4. $-\gamma_0(\theta)$ *is a simple (resp. double) zero of* φ_θ, *if* $\theta>0$ *or* $\theta=0$ *and* $\mathbb{E}(X_1)\neq 0$ *(resp. otherwise, i.e.* $\theta=0$ *and* $\mathbb{E}(X_1)=0$*).*

Proof of Proposition 2.2
1. It is clear that φ_θ is holomorphic in $\{q\in\mathbb{C}\ ;\ \mathrm{Re}\, q>-r_\nu\}$ and the only real zeros of φ_θ in this domain are $-\gamma_0(\theta)$ and $\gamma_0^*(\theta)$.
2. We claim that φ_θ admits only two zeros, $-\gamma_0(\theta)$ and $\gamma_0^*(\theta)$, in $\{q\in\mathbb{C}\ ;\ -\gamma_0(\theta)\leq Req\leq\gamma_0^*(\theta)\}$.

a) Suppose first that $-\gamma_0(\theta)<\mathrm{Re}\, q<\gamma_0^*(\theta)$. We have :

$$|e^{\varphi_\theta(q)}|=|\mathbb{E}(e^{-\theta X_1-\theta})|\leq\mathbb{E}\left(e^{-\mathrm{Re}\,\theta X_1-\theta}\right)=e^{\varphi_\theta(\mathrm{Re}\, q)}<1\,,$$

since $\varphi_\theta<0$ on $]-\gamma_0(\theta),\gamma_0^*(\theta)[$. Then $\varphi_\theta(q)\neq 0$.

b) Let $q = -\gamma_0(\theta) + ib$, $b \in \mathbb{R}$. We compute $\varphi_\theta(q)$:

$$\begin{aligned}\varphi_\theta(q) =& \varphi_\theta(-\gamma_0(\theta)) - \frac{b^2}{2} + \int_{-\infty}^{+\infty} e^{\gamma_0(\theta)y}(cos(by) - 1)\nu(dy) \\ &+ i\left(-b\gamma_0(\theta) + cb - \int_{-\infty}^{+\infty} e^{\gamma_0(\theta)y}\left(sin(by) - by\mathbb{1}_{\{|y|<1\}}\right)\nu(dy)\right).\end{aligned} \tag{2.20}$$

But $\varphi_\theta(-\gamma_0(\theta)) = 0$, then $\mathrm{Re}\,(\varphi_\theta(q)) \leq -\frac{b^2}{2}$. Consequently $\varphi_\theta(q) = 0$ iff $b = 0$.

c) The same reasoning applies to the case $q = -\gamma_0^*(\theta) + ib$.

3. Let us modify the decomposition of φ_θ :

$$\varphi_\theta(q) = \frac{q^2}{2} - cq - \theta + \widehat{\nu_{|[1,+\infty[}}(q) - \nu([1;+\infty[) + \int_{-\infty}^{1} \left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}}\right)\nu(dy)\,. \tag{2.21}$$

Obviously (2.3) implies that $q \to \int_{-\infty}^{1}\left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}}\right)\nu(dy)$ is holomorphic in
$\{q \in \mathbb{C}\ ;\ \mathrm{Re}\,q \leq \beta\}$, for any $\beta \in \mathbb{R}$.

a) Suppose $\theta > 0$ or $\theta = 0$ and $\mathbb{E}(X_1) < 0$. Then φ_θ admits $\gamma_0^*(\theta)$ as a unique positive zero.
The crucial point is the following : assumption (2.16) implies that there exists $R > R_0 > 0$, $k > 0$ such that $|\varphi_\theta(q)| \geq kq^2$ for any q, $-B \leq \mathrm{Re}\,q \leq 0$, $\mathrm{Im}\,q > R$.
Proposition 2.2 will be a direct consequence of the two previous steps 1) and 2), $\varphi_\theta(\overline{z}) = \overline{\varphi_\theta(z)}$ and φ_θ is meromorphic in D_{B_ν}.

b) The case $\theta = 0$ and $\mathbb{E}(X_1) \geq 0$, can be treated similarly. □

Remark 2.3 As it has been pointed out in the previous proof, assumption (2.16) may be replaced by :

$$\exists\varepsilon > 0 \quad \text{such that} \quad \inf_{q\in\mathcal{D}}\left|\frac{\frac{q^2}{2} - cq + \widehat{\nu_{|[1,+\infty[}}(q)}{q^{1+\varepsilon}}\right| > 0 \tag{2.22}$$

where $\mathcal{D} = \{q \in \mathbb{C}\ ;\ -B \leq \mathrm{Re}\,q \leq 0\ ,\ |\mathrm{Im}\,q| \geq R_0\}$.

Theorem 2.4 *We suppose (2.7) and (2.16). Let $B \in]0, B_\nu[$, and $\kappa > 0$, small enough, such that (2.7) holds and (2.19) is satisfied for any $\theta \in [0,\kappa]$ (p being independent from $\theta \in [0,\kappa]$).*

Then for any $\theta \in [0,\kappa]$, $\mu \geq 0$ and $\rho \geq 0$, there exists a positive number $C_0(\theta,\mu,\rho) > 0$ and complex x-polynomial functions $C_1(\theta,\mu,\rho,x),\cdots,C_p(\theta,\mu,\rho,x)$ such that $F(\theta,\mu,\rho,x)$ has the following asymptotic expansion as $x \to +\infty$:

$$\begin{aligned} F(\theta,\mu,\rho,x) =& C_0(\theta,\mu,\rho)e^{-\gamma_0(\theta)x} + \sum_{i=1}^{p} a_i \left(C_i(\theta,\mu,\rho,x)e^{-\gamma_i(\theta)x} + \overline{C_i}(\theta,\mu,\rho,x)e^{-\overline{\gamma_i}(\theta)x} \right) \\ & + \mathrm{O}\left(e^{-Bx}\right) , \end{aligned} \tag{2.23}$$

where $a_i = \dfrac{1}{2}$ if $\gamma_i(\theta)$ is real and $a_i = 1$ otherwise; the degree of $C_i(\theta,\mu,\rho,.)$ is $n_i - 1$, where n_i is the order of multiplicity of $-\gamma_i(\theta)$ and O *is uniform with respect to $\mu \geq 0$, $\rho \geq 0$ and $\theta \in [0,\kappa]$.*

Remark 2.5 *1. If we drop assumption (2.7), then φ_θ has no zero located in $[-r_\nu, 0[$ and the asymptotic expansion (2.23) reduces to :*

$$F(\theta,\mu,\rho,x) = \mathrm{O}(e^{-(r_\nu-\varepsilon)x}) , \quad \varepsilon > 0 . \tag{2.24}$$

2. Heuristically, no assumption on the negative jumps is required to get :

$$F(\theta,\mu,\rho,x) \leq Ce^{-\gamma_0(\theta)x} . \tag{2.25}$$

However to obtain an equivalent, or an asymptotic development of $F(\theta,\mu,\rho,x)$ when x goes to infinity, it is natural to suppose that the negative and the positive parts of the jumps of (X_t) are controlled. Our asymptotic development looks like a perturbation theorem around the case of Brownian motion with negative drift.

3. We give three classes of measures ν satisfying (2.16) :

a) Suppose that ν has finite exponential moments :

$$\forall q \in \mathbb{R} \qquad \int_{-\infty}^{+\infty} \left| e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}} \right| \nu(dy) < +\infty . \tag{2.26}$$

In that case, $\widehat{\nu_{|[1,+\infty[}}$ and φ are holomorphic functions in the whole plane $\mathbb{C}$, then $B_\nu = +\infty$. Moreover, for any $B > 0$:

$$\sup_{\mathrm{Re}\, q \geq -B} \left| \int_1^{+\infty} e^{-qy}\nu(dy) \right| \leq \int_1^{+\infty} e^{By}\nu(dy) < +\infty . \tag{2.27}$$

Then (2.16) holds. Condition (2.26) is realized if ν has a compact support.

b) *Let ν being a linear combination of gamma distributions :*

$$\nu(dy) := \sum_{i=1}^{n} \rho_i e^{-\beta_i y} y^{m_i} \mathbb{1}_{\{y \geq 0\}} dy \,. \tag{2.28}$$

where $\rho_i > 0$, $\beta_i > 0$ and $m_i \in \mathbb{N}$, for any $i \in \{1, 2, \cdots, n\}$,.
Since the Laplace transform of ν is explicit, we obtain immediately its meromorphic extension to the whole plane ($B_\nu = +\infty$) and (2.16). Indeed :

$$\widehat{\nu}(q) = \int_0^{+\infty} e^{-qx} \sum_{i=1}^{n} \rho_i e^{-\beta_i x} x^{m_i} dx = \sum_{i=1}^{n} \rho_i \int_0^{+\infty} x^{m_i} e^{-(q+\beta_i)x} dx \,. \tag{2.29}$$

Setting $y = (q + \beta_i)x$, we have :

$$\widehat{\nu}(q) = \sum_{i=1}^{n} \frac{\rho_i}{(q+\beta_i)^{m_i+1}} \int_0^{+\infty} y^{m_i} e^{-y} dy = \sum_{i=1}^{n} \frac{\rho_i m_i!}{(q+\beta_i)^{m_i+1}} \,. \tag{2.30}$$

This implies that $\widehat{\nu}$ and φ are holomorphic in $\mathbb{C} - \{-\beta_1, \cdots, -\beta_n\}$ and meromorphic on $\mathbb{C}$.

c) *Previous example may be generalized taking :*

$$\nu(dy) := \phi(y) \mathbb{1}_{\{y \geq 0\}} dy \,, \tag{2.31}$$

where $\phi \geq 0$, bounded on $[0, y_0]$, and for every $y \geq y_0$:

$$\phi(y) := \rho_0 e^{-\beta_0 y} y^{m_0 - 1} + \sum_{i=1}^{n} (\rho_i e^{-\beta_i y} + \overline{\rho_i} e^{-\overline{\beta_i} y}) y^{m_i - 1} + \mathrm{O}(e^{-\beta_{n+1}\, y}) \,, \tag{2.32}$$

with $y_0 \geq 0$, $\rho_0 \geq 0$, $\beta_0 > 0$, $\mathrm{Re}\,(\beta_i) > 0$, $\rho_i \in \mathbb{C}^*$, $m_i \in \mathbb{N}^*$ et $\beta_{n+1} \geq \sup_{i \in \{1, \cdots, n\}} \mathrm{Re}\,(\beta_i)$.
Then $\widehat{\nu}$ and φ are meromorphic functions on $\{q \,/\, \mathrm{Re}\, q \geq \beta_{n+1}\}$, $B_\nu = \beta_{n+1}$ and (2.16) holds.

d) *If $\nu = \nu_1 + \nu_2$, ν_1 and ν_2 verify (2.16) then ν also.*

Proof of Theorem 2.4
Recall (cf. Theorem 2.1 of [12]) that $\widehat{F}(\theta, \mu, \rho, .)$ verifies :

$$\widehat{F}(\theta, \mu, \rho, q) = \frac{1}{\varphi(q) - \theta} \left(\frac{q - \gamma_0^*(\theta)}{2} + \int_0^{+\infty} \left[\frac{e^{-(q+\rho)y} - e^{-\mu y}}{q + \rho - \mu} - \frac{e^{-(\gamma_0^*(\theta)+\rho)y} - e^{-\mu y}}{\gamma_0^*(\theta) + \rho - \mu} \right] \nu(dy) \right.$$
$$\left. + RF(\theta, \mu, \rho, .)(q) - RF(\theta, \mu, \rho, .)(\gamma_0^*(\theta)) \right) \,, \tag{2.33}$$

where $\operatorname{Re} q > 0$ and R denotes the operator :

$$Rh(q) := \int_{-\infty}^{0} \nu(dy) \int_{0}^{-y} \left(e^{-q(b+y)} - 1\right) h(b)db \,. \tag{2.34}$$

For simplicity we only consider the case $\rho = 0$. The function $F(\theta, \mu, 0, \cdot)$ will be written $F(\theta, \mu, \cdot)$ for short.
In the sequel, $B \in]0,\, B_\nu[$ is supposed to be close to B_ν.

Step 1 : Replacing $F(\theta, \mu, .)$ by $\widetilde{F}(\theta, \mu, .)$ and equation associated with $\widetilde{F}(\theta, \mu, .)$
We entend continuously $F(\theta, \mu, .)$ to the whole line in the following manner :

$$\widetilde{F}(\theta, \mu, x) := F(\theta, \mu, x)\mathbb{1}_{[0;+\infty[}(x) + (1+x)\mathbb{1}_{[-1;+0]}(x), \qquad \forall x \in \mathbb{R}\,. \tag{2.35}$$

Let $\widehat{\widetilde{F}}(\theta, \mu .)$ be the Laplace transform of $\widetilde{F}(\theta, \mu, .)$:

$$\widehat{\widetilde{F}}(\theta, \mu, q) := \int_{-\infty}^{+\infty} e^{-qx} \widetilde{F}(\theta, \mu, x)dx \,. \tag{2.36}$$

The advantage of using $\widetilde{F}(\theta, \mu, .)$ instead of $F(\theta, \mu, .)$ lies in the fact that we shall prove in Lemma 2.6 below, that $t \to \widehat{\widetilde{F}}(\theta, \mu, q_1 + it)$ is an integrable function on $\mathbb{R}$, for any q_1 in $]0, \beta_\theta[$ (β_θ being defined in Proposition 2.2).
Since $\displaystyle\int_{-1}^{0} (1+x)e^{-qx}dx = \frac{e^q - 1 - q}{q^2}$, (2.33) implies :

$$\begin{aligned}
\widehat{\widetilde{F}}(\theta, \mu, q) =& \frac{e^q - 1 - q}{q^2} \\
&+ \frac{1}{\varphi(q) - \theta} \Bigg(\frac{q - \gamma_0^*(\theta)}{2} + \int_{0}^{+\infty} \left[\frac{e^{-qy} - e^{-\mu y}}{q - \mu} - \frac{e^{-\gamma_0^*(\theta) y} - e^{-\mu y}}{\gamma_0^*(\theta) - \mu} \right] \nu(dy) \\
&+ RF(\theta, \mu, .)(q) - RF(\theta, \mu, .)(\gamma_0^*(\theta)) \Bigg)
\end{aligned} \tag{2.37}$$

We observe that

$$\begin{aligned}
&\frac{\varphi(q) - \theta}{q + \gamma_0^*(\theta)} = \frac{\varphi(q) - \varphi(\gamma_0^*(\theta))}{q + \gamma_0^*(\theta)} \\
&= \frac{q - \gamma_0^*(\theta)}{2} + c\, \frac{q - \gamma_0^*(\theta)}{q + \gamma_0^*(\theta)} + \int_{-\infty}^{+\infty} \frac{e^{-qy} - e^{-\gamma_0^*(\theta) y} + (q - \gamma_0^*(\theta)) y \mathbb{1}_{\{|y|<1\}}}{q + \gamma_0^*(\theta)} \nu(dy) \,,
\end{aligned} \tag{2.38}$$

and

$$\frac{e^q - 1 - q}{q^2} + \frac{1}{q + \gamma_0^*(\theta)} = \frac{e^q - 1}{q^2} - \frac{\gamma_0^*(\theta)}{q(q + \gamma_0^*(\theta))} \,. \tag{2.39}$$

Consequently for any q, such that $\operatorname{Re} q > 0$:

$$
\begin{aligned}
\widehat{\widehat{F}}(\theta,\mu,q) = {} & \frac{e^q-1}{q^2} - \frac{\gamma_0^*(\theta)}{q(q+\gamma_0^*(\theta))} \\
& + \frac{1}{\varphi(q)-\theta}\left[-c\,\frac{q-\gamma_0^*(\theta)}{q+\gamma_0^*(\theta)} - \int_{-\infty}^{+\infty} \frac{e^{-qy}-e^{-\gamma_0^*(\theta)y}+(q-\gamma_0^*(\theta))y\mathbb{1}_{\{|y|<1\}}}{q+\gamma_0^*(\theta)}\nu(dy) \right. \\
& + \int_0^{+\infty}\left[\frac{e^{-qy}-e^{-\mu y}}{q-\mu}\nu(dy) - \frac{e^{-\gamma_0^*(\theta)y}-e^{-\mu y}}{\gamma_0^*(\theta)-\mu}\right]\nu(dy) \\
& \left. + RF(\theta,\mu,.)(q) - RF(\theta,\mu,.)(\gamma_0^*(\theta)) \right]. \qquad (2.40)
\end{aligned}
$$

It is clear that (2.3) implies that $RF(\theta,\mu,.)$ is an entire function on $\mathbb{C}$ and all the integrals in (2.40) are holomorphic at least in D_{B_ν}. Consequently the right hand-side of (2.40) is the meromorphic extension of $\widehat{\widehat{F}}(\theta,\mu,.)$ to D_{B_ν}.

Before ending step 1, we remark that if $\theta > 0$ or $\theta = 0$ and $\mathbb{E}(X_1) \neq 0$, $\widehat{\widehat{F}}(\theta,\mu,.)$ and $\widehat{F}(\theta,\mu,.)$ are holomorphic in a neighbourhood of $q = \gamma_0^*(\theta)$, although $\gamma_0^*(\theta)$ is a zero of φ_θ, this value being a false singularity for $\widehat{\widehat{F}}(\theta,\mu,.)$. But if $\theta = 0$ and $\mathbb{E}(X_1) = 0$, then $\gamma_0^*(0) = 0$ is a pole for $\widehat{\widehat{F}}(0,\mu,.)$.

Step 2 : $t \to \widehat{\widehat{F}}(\theta,\mu,q_1+it)$ **belongs to** $\mathbb{L}^1(\mathbb{R})$

Let θ be a fixed element in $[0,\kappa]$, $\beta_\theta > 0$, given by Proposition 2.2 and $q_1 \in]0,\beta_\theta[$.

Lemma 2.6 *Under (2.7) and (2.16), the function* $t \to \widehat{\widehat{F}}(\theta,\mu,q_1+it)$ *belongs to* $\mathbb{L}^1(\mathbb{R})$.

To prove Lemma 2.6, we begin with stating few technical inequalities. These relations will be also used in the sequel.

Lemma 2.7 *Let $\theta_1 < \theta_2$. We suppose (2.16) is realized. Then :*

(i) $\quad \exists k > 0$ such that for any q satisfying $\operatorname{Re} q \in [\theta_1, \theta_2]$, we have :

$$\left| \int_{-1}^{1} \left(e^{-qy} - 1 + qy \mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| \le k|q| \, , \tag{2.41}$$

$$\left| \int_{-1}^{0} \left(e^{-qy} - 1 + qy \mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| \le k|q| , \tag{2.42}$$

$$\left| \int_{0}^{1} \left(e^{-qy} - 1 + qy \mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| \le k|q| \, , \tag{2.43}$$

(ii) $\quad \forall d > 0, \quad \exists k_0 > 0$ such that $\displaystyle \sup_{\operatorname{Re} q \le d} \left| \int_{-\infty}^{-1} e^{-qy} \nu(dy) \right| \le k_0 \qquad (2.44)$

(iii) $\quad \forall A > 0 \quad \exists k_1 > 0$ such that $\forall q$ satisfying : $\left\{ \begin{array}{c} -B \le \operatorname{Re} q \le A \\ |\operatorname{Im} q| \ge R_0 \end{array} \right.$,

$$\left| \int_{-\infty}^{+\infty} \left(e^{-qy} - 1 + qy \mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| \ \le \ k_1 (1 + |q|) \, , \tag{2.45}$$

$$\frac{1}{k_1} \, |q|^2 \ \le \ |\varphi(q)| \ \le \ k_1 \, |q|^2 \tag{2.46}$$

(iv) $\quad \forall h \in \mathbb{R}, \quad \displaystyle \sup_{\operatorname{Re} q \le h} |RF(\theta, \mu, q)| < +\infty \, . \qquad (2.47)$

Proof of Lemma 2.7

(i) We set $q = a + ib$, where $a \in [\theta_1, \theta_2]$. We have :

$$\left| \int_{-1}^{1} \left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| \leq \left| \int_{-1}^{1} \left(e^{-(a+ib)y} - e^{-ay} + iby\mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| + \left| \int_{-1}^{1} \left(e^{-ay} - 1 + ay\mathbb{1}_{\{|y|<1\}} \right) \nu(dy) \right| . \tag{2.48}$$

We deduce immediately (2.41).
The two other inequalities (2.42) and (2.43) can be proved by the same way.

(ii) Let q such that $\operatorname{Re} q \leq d$, then

$$\left| \int_{-\infty}^{-1} e^{-qy} \nu(dy) \right| \leq \int_{-\infty}^{-1} e^{-\operatorname{Re} qy} \nu(dy) \leq \int_{-\infty}^{-1} e^{-dy} \nu(dy) = k_0 \, . \tag{2.49}$$

Assumption (2.3) implies that k_0 is a finite constant.

(iii) For any q satisfying $-B \leq \operatorname{Re} q \leq A$ and $|\operatorname{Im} q| \geq R_0$, we get

$$\int_{-\infty}^{+\infty} \left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}} \right) \nu(dy) = \int_{-\infty}^{-1} e^{-qy} \nu(dy) + \int_{-1}^{1} \left(e^{-qy} - 1 + qy\mathbb{1}_{\{|y|<1\}} \right) \nu(dy) + \int_{1}^{+\infty} e^{-qy} \nu(dy) + \int_{-\infty}^{-1} \nu(dy) + \int_{1}^{+\infty} \nu(dy) \, . \tag{2.50}$$

Relation (2.45) follows from (2.16), (2.41) and (2.42).

It is easy to check (2.46) by virtue of the definition of φ (cf. (2.1)) and previous inequalities.

(iv) Using the definition of RF (cf. (2.34)), we deduce that $q \to RF(\theta, \mu, q)$ is an increasing function on $\mathbb{R}$ vanishing at 0.
If q is a complex number such that $\operatorname{Re} q \leq h$, then

$$\begin{aligned} |RF(\theta, \mu, q)| &\leq \int_{-\infty}^{0} \left(\int_{0}^{-y} e^{-\operatorname{Re} q(y+b)} F(\theta, \mu, b) db - y \right) \nu(dy) \\ &\leq k \; RF(\theta, \mu, h) \end{aligned}$$

This implies (2.47). □

Proof of Lemma 2.6

Proposition 2.2 tells us φ_θ has no zero in the strip $\{q \in \mathbb{C} \;/\; 0 < \operatorname{Re} q < \beta_\theta\}$. Hence if $q_1 \in]0, \beta_\theta[$, $\varphi(q_1+it)-\theta$ neither cancels and (2.40) implies that $t \to \widehat{\widetilde{F}}(\theta, \mu, q_1 + it)$ is a continuous function. Let us focus on (2.40). Lemma 2.7 implies that all the numerators in (2.40) are bounded on the line $\{q_1 + it \;/\; t \in \mathbb{R}\}$, and the denominators are less than $C|q^2|$ when $|q| \to +\infty$ where $C > 0$. This proves that $t \to \widehat{\widetilde{F}}(\theta, \mu, q_1 + it)$ belongs to $\mathbb{L}^1(\mathbb{R})$. $\square$

Step 3 : Proof of the asymptotic development (2.23), through the Mellin Fourier inverse transform

Proposition 2.2 gives the existence of $\kappa > 0$ and B such that for any $\theta \in [0, \kappa]$, φ_θ does not vanish on $\{-B + it \;/\; t \in \mathbb{R}\}$ and $B \neq \gamma_0^*(\theta)$.

Let $0 < q_1 < \beta_\theta$. Since $t \to \widehat{\widetilde{F}}(\theta, \mu, q_1 + it)$ belongs to $\mathbb{L}^1(\mathbb{R})$ (cf. Lemma 2.6), we are allowed to make use of the Mellin Fourier inverse transform. So, for any $x \geq 0$:

$$e^{-q_1 x}\widetilde{F}(\theta, \mu, x) = e^{-q_1 x} F(\theta, \mu, x) = \frac{1}{2\pi}\int_{-\infty}^{+\infty} e^{itx}\widehat{\widetilde{F}}(\theta, \mu, q_1 + it)dt\,, \tag{2.51}$$

hence

$$F(\theta, \mu, x) = \frac{1}{2\pi}\int_{-\infty}^{+\infty} e^{(q_1+it)x}\widehat{\widetilde{F}}(\theta, \mu, q_1 + it)dt = -\frac{i}{2\pi}\int_{\Gamma_{q_1}} e^{zx}\widehat{\widetilde{F}}(\theta, \mu, z)dz\,, \tag{2.52}$$

where Γ_{q_1} is the path :

$$\Gamma_{q_1} := \{z = q_1 + it \quad \text{such that} \quad t \in \mathbb{R}, \quad t \text{ increasing}\}\,. \tag{2.53}$$

In Proposition 2.2, it is proved there exists $R_1 > R_0$, such that φ_θ has no zero in the two half-strips $\{q \in \mathbb{C} \;/ -B \leq \operatorname{Re} q < \beta_\theta \text{ and } |\operatorname{Im} q| > R_1\}$. In particular $\widehat{\widetilde{F}}(\theta, \mu, .)$ is holomorphic in this domain.

Let $\Gamma_{-B,q_1,R}$ be the rectangular path (see Figure 1) :

$$\Gamma_{-B,q_1,R} := \Gamma_{q_1,R} \,\cup\, \Gamma_R \,;\cup\, \Gamma_{-B,R} \,\cup\, \Gamma_{-R}\,, \tag{2.54}$$

where :

$$\Gamma_{q_1,R} := \{q_1 + it \;/ \quad |t| \leq R, \quad t \text{ growing}\}\,, \tag{2.55}$$

$$\Gamma_R := \{t + iR \;/ \quad -B \leq t \leq q_1, \quad t \text{ decreasing}\} \tag{2.56}$$

$$\Gamma_{-B,R} := \{-B + it \;/ \quad |t| \leq R, \quad t \text{ decreasing}\}\,, \tag{2.57}$$

$$\Gamma_{-R} := \{t - iR \;/ \quad -B \leq t \leq q_1, \quad t \text{ growing}\}\,. \tag{2.58}$$



Figure 1: The path $\Gamma_{-B,q_1,R}$

Applying the residual theorem to the meromorphic extension of $z \to e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)$ to D_{B_ν}, we obtain, for any $R > R_1$:

$$\int_{\Gamma_{-B,q_1,R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz =$$

$$2i\pi\left[C_0(\theta,\mu)e^{-\gamma_0(\theta)x} + \sum_{i=1}^{p} a_i\left(C_i(\theta,\mu,x)e^{-\gamma_i(\theta)x} + \overline{C_i}(\theta,\mu,x)e^{-\overline{\gamma_i}(\theta))x}\right)\right] \qquad (2.59)$$

where $a_i = \dfrac{1}{2}$ if $-\gamma_i(\theta)$ is a real number and $a_i = 1$ otherwise, and :

$$C_0(\theta,\mu) := Res\left(\widehat{\widehat{F}}(\theta,\mu,z);\ -\gamma_0(\theta)\right)\ , \qquad (2.60)$$

$$C_i(\theta,\mu,x) := e^{\gamma_i(\theta)x}Res\left(e^{zx}\widehat{\widehat{F}}(\theta,\mu,z);\ -\gamma_i(\theta)\right)\ , \qquad (2.61)$$

where $Res(f(z);\gamma)$ denotes the residual of f at point γ. Let us remark (2.60) is valid since $-\gamma_0(\theta)$ is a simple pole of $\widehat{\widehat{F}}(\theta,\mu,.)$.

Since $z \to \dfrac{e^z-1-z}{z^2}$ has an holomorphic extension to the whole plan $\mathbb{C}$, identity

(2.35) implies that :

$$C_0(\theta,\mu) = Res\left(\widehat{F}(\theta,\mu,z);\ -\gamma_0(\theta)\right)\,, \tag{2.62}$$

$$C_i(\theta,\mu,x) = e^{\gamma_i(\theta)} Res\left(e^{zx}\widehat{F}(\theta,\mu,z);\ -\gamma_i(\theta)\right) \qquad \forall i \in \{1,\cdots,p\}\,. \tag{2.63}$$

We observe that $\overline{C_i}(\theta,\mu,x) = e^{\overline{\gamma_i}(\theta)x} Res\left(e^{zx}\widehat{F}(\theta,\mu,z);\ -\overline{\gamma_i}(\theta)\right)$. We will prove in Remark 2.8 below that $x \to C_i(\theta,\mu,x)$ is a polynomial function.
Since $z \to e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)$ belongs to $\mathbb{L}^1(\mathbb{R})$ (cf. (2.51)), we have :

$$\begin{aligned} F(\theta,\mu,x) &= -\frac{i}{2\pi}\lim_{R\to+\infty}\int_{\Gamma_{q_1,R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz \\ &= -\frac{i}{2\pi}\lim_{R\to+\infty}\left[\int_{\Gamma_{-B,q_1,R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz - \int_{\Gamma_R} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz\right. \\ &\qquad\qquad \left. - \int_{\Gamma_{-B,R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz - \int_{\Gamma_{-R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz\right]. \end{aligned} \tag{2.64}$$

We claim that in the right hand-side of (2.64) the limits of the second term and the fourth one are null. As for the third limit, we have :

$$\lim_{R\to+\infty}\int_{\Gamma_{-B,R}} e^{zx}\widehat{\widehat{F}}(\theta,\mu,z)dz = \mathrm{O}(e^{-Bx})\,, \tag{2.65}$$

where O is uniform with respect to $\theta \in [0,\kappa]$ and $\mu \in \mathbb{R}_+$.
Hence, as $x \to +\infty$:

$$F(\theta,\mu,x) = C_0(\theta,\mu)e^{-\gamma_0(\theta)} + \sum_{i=1}^{p} a_i\Big[C_i(\theta,\mu,x)e^{-\gamma_i(\theta)x} + \overline{C_i}(\theta,\mu,x)e^{-\overline{\gamma_i}(\theta)x}\Big] + \mathrm{O}(e^{-Bx})\,. \tag{2.66}$$

Let us determine the sign of $C_0(\theta,\mu)$. The asymptotic expansion (2.23) implies that :

$$\lim_{x\to+\infty} e^{\gamma_0(\theta)x}F(\theta,\mu,x) = C_0(\theta,\mu) \geq 0\,. \tag{2.67}$$

The residual $C_0(\theta,\mu)$ at $-\gamma_0(\theta)$ of $F(\theta,\mu,.)$ cannot vanish because $-\gamma_0(\theta)$ is a single pole of $\widehat{F}(\theta,\mu,.)$ when $\theta > 0$ or $\theta = 0$ and $\mathbb{E}(X_1) \neq 0$.
If $\theta = 0$ and $c = \mathbb{E}(J_1)$, it is easy to see that $C_0(0,\mu) \neq 0$ (cf. (2.73)). □

Remark 2.8 *1. We would like to show that $C_i(\theta,\mu,\rho,.)$ is a polynomial function and determine its maximal degree. Assume that $-\gamma_i(\theta)$ is a zero of φ_θ with* *multiplicity n_i, and $\widehat{F}(\theta,\mu,\rho,z)$ has the following asymptotic expansion in a neighborhood of $-\gamma_i(\theta)$:*

$$\widehat{F}(\theta,\mu,\rho,z)=\frac{K_{i,n_i}(\theta,\mu,\rho)}{(z+\gamma_i(\theta))^{n_i}}+\frac{K_{i,n_i-1}(\theta,\mu,\rho)}{(z+\gamma_i(\theta))^{n_i-1}}+\cdots+\frac{K_{i,1}(\theta,\mu,\rho)}{z+\gamma_i(\theta)}+\cdots\,. \tag{2.68}$$

Since

$$e^{zx}=e^{-\gamma_i(\theta)x}\left(1+(z+\gamma_i(\theta))x+\frac{(z+\gamma_i(\theta))^2}{2!}x^2+\cdots\right) \tag{2.69}$$

then

$$C_i(\theta,\mu,\rho,x)=\frac{K_{i,n_i}(\theta,\mu,\rho)}{(n_i-1)!}x^{n_i-1}+\frac{K_{i,n_i-1}(\theta,\mu,\rho)}{(n_i-2)!}x^{n_i-2}+\cdots+K_{i,1}(\theta,\mu,\rho)\,. \tag{2.70}$$

2. *(i) Suppose that $-\gamma_i(\theta)$ is a single zero of φ_θ (it is also a single pole of $\widehat{F}(\theta,\mu,.)$), then $C_i(\theta,\mu,\rho,x)$ does not depend on x and is given by the following :*

$$C_i(\theta,\mu,\rho)=Res\left(\widehat{F}(\theta,\mu,\rho,z);\ -\gamma_i(\theta)\right)\,. \tag{2.71}$$

This situation occurs if $i=0$, when $\theta>0$ or $\theta=0$ and $\mathbb{E}(X_1)\neq 0$.
If the real part of $-\gamma_i(\theta)$ is bigger than $-r_\nu$, $C_i(\theta,\mu,\rho)$ can be determined as follows :

$$\begin{aligned}C_i(\theta,\mu,\rho)=&\frac{1}{\varphi'(-\gamma_i(\theta))}\left[\frac{-\gamma_i(\theta)-\gamma_0^*(\theta)}{2}\right.\\&+\int_0^{+\infty}\left[\frac{e^{(\gamma_i(\theta)-\rho)y}-e^{-\mu y}}{-\gamma_i(\theta)+\rho-\mu}-\frac{e^{-(\gamma_0^*(\theta)+\rho)y}-e^{-\mu y}}{\gamma_0^*(\theta)+\rho-\mu}\right]\nu(dy)\\&\left.+RF(\theta,\mu,\rho,.)(-\gamma_i(\theta))-RF(\theta,\mu,\rho,.)(\gamma_0^*(\theta))\right]\,,\end{aligned} \tag{2.72}$$

where it is supposed that $\dfrac{e^{ay}-1}{a}=y$ if $a=0$.
When $\mathrm{Re}\,(-\gamma_i(\theta))<-r_\nu$, the previous ν-integrals and φ, have to be replaced by their meromorphic extensions.
Recall that if $\mathbb{E}(X_1)<0$ (resp. $\mathbb{E}(X_1)>0$) then $\gamma_0^*(0)=0$ (resp. $\gamma_0(0)=0$).

(ii) If $\mathbb{E}(X_1) = 0$, then $\gamma_0(0) = 0$ is a double zero of φ, but a simple pole of $\widehat{F}(\theta, \mu, \rho, .)$. Thus, by a direct calculation, we have :

$$C_0(0, \mu, \rho) = \frac{1}{\varphi''(0)} \left(1 - \frac{2}{(\rho - \mu)^2} \int_0^{+\infty} e^{-\rho y} \left(1 - e^{(\rho - \mu)y} + (\rho - \mu)y\right) \nu(dy) \right.$$
$$\left. -2 \int_{-\infty}^0 \nu(dy) \int_0^{-y} (y + b) F(0, \mu, \rho, b) db \right) . \qquad (2.73)$$

In particular :

$$C_0(0, 0, 0) = \frac{1}{\varphi''(0)} \left(1 + \int_0^{+\infty} y^2 \, \nu(dy) - 2 \int_{-\infty}^0 \nu(dy) \int_0^{-y} (y + b) F(0, 0, b) db\right)$$
$$= 1 - \frac{2}{\varphi''(0)} \int_{-\infty}^0 \nu(dy) \int_0^{-y} (y + b) F(0, 0, 0, b) db \, . \qquad (2.74)$$

3. *In [1], it was proved that $\mathbb{P}(T_x < +\infty) \sim C_0 e^{-\gamma_0 x}$ as $x \to +\infty$, C_0 being indeterminate. We go a little bit further since $C_0 = C_0(0, 0, 0)$. If we suppose moreover that the support of ν is included in $[0, +\infty[$, then $RF(\theta, \mu, \rho, .) = 0$ and*

$$C_i(0, 0, 0) = -\frac{\varphi'(0)}{\varphi'(-\gamma_i(0))}, \quad \text{if } \mathbb{E}(X_1) < 0 \, , \qquad (2.75)$$
$$C_i(0, 0, 0) = 0, \ i \neq 0 \ \text{ and } \ C_0(0, 0, 0) = 1, \quad \text{if } \mathbb{E}(X_1) \geq 0 \, . \qquad (2.76)$$

We recover the result given in [2], i.e. (2.75) with $i = 0$.

3 A polynomial upper bound for the ruin probability

In this section we only consider the ruin probability. For simplicity we note :

$$F(x) := F(0, 0, 0, x) = \mathbb{P}(T_x < +\infty) \, , \qquad (3.1)$$

and its Laplace transform :

$$\widehat{F}(q) := \int_0^{+\infty} e^{-qx} F(x) dx \qquad \forall q \in \mathbb{C} \, , \ \mathrm{Re}\,(q) > 0 \, . \qquad (3.2)$$

We suppose in this section :

$$\int_{|y|>1} |y| \nu(dy) < +\infty \qquad \text{and} \qquad \mathbb{E}(X_1) = \int_{|y|>1} y \nu(dy) - c < 0 \, . \qquad (3.3)$$

Therefore $\lim\limits_{x \to +\infty} \mathbb{P}(T_x < +\infty) = 0$.
In Section 2 we have proved that, under suitable assumptions, the ruin probability goes to 0, with exponential rate. Here the aim is to prove that under suitable assumptions, $F(x)$ has a polynomial type rate of decay, as $x \to \infty$.

Theorem 3.1 *We suppose (3.3) and*

$$\int_0^{+\infty} y^p \nu(dy) < +\infty \,, \quad \text{for some} \quad p \geq 2\,. \tag{3.4}$$

Let n be the integer part of $p-2$, then

$$\forall x \in \mathbb{R}_+ \qquad F(x) \leq \frac{C_n}{1+x^n}\,, \tag{3.5}$$

where $C_n > 0$.

Proof of Theorem 3.1
The proof will be divided into five parts.

Step 1 We prove that it is sufficient to consider a Lévy measure ν with support included in $[-k, +\infty[$, for some (finite) $k \geq 0$.

Assumption (3.3) implies there exists $k > 0$ such that

$$\int_{-k}^{+\infty} \mathbb{1}_{\{|y|>1\}} y \nu(dy) < c\,. \tag{3.6}$$

Let $(X_t^k\,,\ t \geq 0)$ be a Lévy process with decomposition :

$$X_t^k := B_t - ct + J_t^k \qquad \forall t \geq 0\,, \tag{3.7}$$

where $(J_t^k\,,\ t \geq 0)$ is a pure jump process with Lévy measure $\nu_k = \nu_{|_{[-k;+\infty[}}$, independent of $(B_t\,,\ t \geq 0)$. Moreover $(J_t^k\,,\ t \geq 0)$, $(J_t\,,\ t \geq 0)$ can be defined on the same probability space, and $J_t \leq J_t^k$, $\forall t \geq 0$. Then a.e. $X_t \leq X_t^k$, $\forall t \geq 0$ and $T_x^k \leq T_x$, $\forall x \geq 0$ where

$$T_x^k := \inf\{t \geq 0 \ / \ X_t^k > x\}\,. \tag{3.8}$$

As a result

$$\forall x \geq 0 \qquad F(x) \leq F^k(x) := \mathbb{P}(T_x^k < +\infty)\,. \tag{3.9}$$

This proves the claim.

In the sequel we suppose that the support of ν is included in $[-k, +\infty[$, for some $k > 0$.

Step 2 F belongs to $\mathbb{L}^1(\mathbb{R}_+)$.

We first prove :

$$\sup_{0<q\le q_0} |\widehat{F}(q)| < +\infty\,, \quad \text{for some } q_0>0\,. \tag{3.10}$$

Choosing $\theta=\mu=\rho=0$ in (2.33), we obtain :

$$\widehat{F}(q)=\frac{1}{\varphi(q)}\left(\frac{q}{2}+\frac{1}{q}\int_0^{+\infty}(e^{-qy}-1+qy)\nu(dy)+RF(q)\right)\,. \tag{3.11}$$

Let us determine the asymptotic behaviour of the numerator and the denominator, $q\to 0$, we have :

$$\varphi(q)\sim_{q\to 0} q\varphi'(0)=q\left(c-\int_{|y|>1}y\nu(dy)\right)\,, \tag{3.12}$$

$$\frac{q}{2}+\frac{1}{q}\int_0^{+\infty}(e^{-qy}-1+qy)\nu(dy)\sim_{q\to 0}\frac{q}{2}\left[1+\int_0^{+\infty}y^2\nu(dy)\right]\,. \tag{3.13}$$

Since the support of ν is included in $[-k,+\infty[$, $RF(q)$ can be simplified :

$$RF(q)=\int_{-k}^{0}e^{-qy}\nu(dy)\left[\int_0^{-y}e^{-qb}F(b)db-\int_0^{-y}F(b)db\right]\,. \tag{3.14}$$

But $RF(0)=0$ and the derivative of $RF(q)$ is bounded, then $|RF(q)|\le Cq$, for any $0\le q\le q_0$. (3.10) follows immediately.

It is now easy to check that F is in $\mathbb{L}^1(\mathbb{R}_+)$. F being positive :

$$\int_0^{+\infty}F(x)dx=\lim_{q\to 0}\int_0^{+\infty}e^{-qx}F(x)dx\le\sup_{0<q\le q_0}\widehat{F}(q)<+\infty\,. \tag{3.15}$$

The function F can be extended to the whole line, setting $F(x)=0$, for any $x\le 0$. However F may have a jump at 0. Let $\widetilde{F}$ be the following continuous extension of F :

$$\widetilde{F}(x):=F(x)\mathbb{1}_{[0;+\infty[}(x)+(1+x)\mathbb{1}_{[-1;+0]}(x),\qquad \forall x\in\mathbb{R}\,. \tag{3.16}$$

Let $q\to\widehat{\widetilde{F}}(iq)$ be the Fourier transform of $\widetilde{F}$:

$$\widehat{\widetilde{F}}(iq)=\int_{-\infty}^{+\infty}e^{-iqx}\widetilde{F}(x)dx=\int_{-1}^{+\infty}e^{-iqx}\widetilde{F}(x)dx\qquad\forall q\in\mathbb{R}\,. \tag{3.17}$$

Step 3 $q \to \widehat{\widetilde{F}}(iq)$ is in $\mathbb{L}^1(\mathbb{R})$.

For simplicity, we suppose moreover that $\displaystyle\int_{-1}^{1} |y|\nu(dy) < +\infty$.

Since $F \in \mathbb{L}^1(\mathbb{R}_+)$, then $\widetilde{F} \in \mathbb{L}^1(\mathbb{R})$ and $\widehat{\widetilde{F}}(i.)$ is continuous.
Consequently if we establish :

$$\left|\widehat{\widetilde{F}}(iq)\right| \le \frac{C}{1+q^2}\,, \tag{3.18}$$

then $\widehat{\widetilde{F}}$ will be an element of $\mathbb{L}^1(\mathbb{R})$.
It is proved in [12] (Annex, Proposition A-2), that $\varphi(iq) = 0$, $q \in \mathbb{R}$ iff $q = 0$.
Therefore we are allowed to replace q by iq in (3.11).
Using the identity :

$$\int_{-1}^{0} (1+x)e^{-iqx}dx = -\frac{e^{iq} - 1 - iq}{q^2}\,, \tag{3.19}$$

we then deduce :

$$\widehat{\widetilde{F}}(iq) = \frac{1-e^{iq}}{q^2} + \frac{1}{\varphi(iq)}\left[-c + \int_{0}^{+\infty} y\nu(dy) + \frac{i}{q}\int_{-k}^{0}(e^{-iqy}-1)\nu(dy) + RF(iq)\right]. \tag{3.20}$$

Since $\varphi(iq) \sim_{|q|\to+\infty} -\dfrac{q^2}{2}$, (3.18) is a consequence of the three inequalities :

$$\left|\frac{1-e^{iq}}{q^2}\right| \le \frac{2}{q^2}\,, \tag{3.21}$$

$$\left|\frac{i}{q}\int_{-k}^{0}(e^{-iqy}-1)\nu(dy)\right| \le \int_{-k}^{0}|y|\nu(dy)\,, \tag{3.22}$$

$$|RF(iq)| \le 2\int_{-k}^{0}|y|\nu(dy)\,. \tag{3.23}$$

Step 4 The $n-2$ first derivatives of $\widehat{\widetilde{F}}$ belong to $\mathbb{L}^1(\mathbb{R})$.

Obviously any k derivative of $q \to \dfrac{1-e^{iq}}{q^2}$ is continuous, bounded by $\dfrac{C}{q^2}$, $|q| \ge 1$, and consequently belongs to $\mathbb{L}^1(\mathbb{R})$.
The second term in the right hand-side of (3.20) may be written as $\dfrac{N(q)}{\varphi(iq)}$.
Assumption (3.4) implies that the n first derivatives of N are bounded, hence $\left|\dfrac{N(q)}{\varphi(iq)}\right| \le \dfrac{C}{q^2}$, $|q| \ge 1$.
As for the asymptotic behaviour of $\dfrac{N(q)}{\varphi(iq)}$ in a neighborhood of 0, it can be proved by a similar reasoning that this ratio is bounded, for any $|q| \le 1$.

Step 5 Proof of (3.5).

Since the $n-2$ derivatives of $q \to \widehat{\widetilde{F}}(iq)$ belong to $\mathbb{L}^1(\mathbb{R})$, then

$$x^{n-2}\widetilde{F}(x) = \frac{i^{n-2}}{2\pi} \int_{\mathbb{R}} e^{iqx} \frac{d^{n-2}}{dq^{n-2}} \left(\widehat{\widetilde{F}}(iq) \right) dq \,. \tag{3.24}$$

This identity directly implies (3.5). □

4 Normalized limit distribution of (T_x, K_x, L_x), as $x \to +\infty$

In this section we investigate the limit behaviour of (T_x, K_x, L_x), as $x \to +\infty$. Recall that K_x and L_x are defined by (1.6). Since $x \to T_x$ is non-decreasing, T_x needs to be normalized. We have to consider three cases either $\mathbb{E}(X_1) > 0$, or $\mathbb{E}(X_1) < 0$ or $\mathbb{E}(X_1) = 0$. Let us start with the case $\mathbb{E}(X_1) < 0$.

Theorem 4.1 *Under hypothesis of Theorem 2.4 and* $\mathbb{E}(X_1) < 0$*, then, conditionally on* $\{T_x < +\infty\}$*,* $\left(\frac{1}{\sqrt{x}} \left(T_x + \frac{x}{\varphi'(-\gamma_0(0))} \right), K_x, L_x \right)$ *converges in distribution to the 3-dimensional law*
$\mathcal{N}\left(0; -\frac{\varphi''(-\gamma_0(0))}{\varphi'^3(-\gamma_0(0))}\right) \otimes w$ *where* w *is the probability measure on* $\mathbb{R}_+ \times \mathbb{R}_+$ *:*

$$\begin{aligned} w(dk, dl) =& \frac{-1}{\mathbb{E}(X_1)} \left[\frac{\gamma_0(0)}{2}\, \delta_{0,0}(dk, dl) + (e^{\gamma_0(0)l} - 1) \mathbb{1}_{\{k \geq 0; l \geq 0\}}\, \nu_l(dk)\, dl \right. \\ & \left. + \int_{\mathbb{R}_-} \left(\int_0^{-y} (1 - e^{\gamma_0(0)(b+y)}) F(0,0,0,b) n(b, dk, dl) db \; \mathbb{1}_{\{k \geq 0, l \geq 0\}} \right) \nu(dy) \right], \end{aligned} \tag{4.1}$$

$\nu_l(dk)$ is the image of $\nu(dk)$ by the map $y \to y - l$, and $n(b, dk, dl)$ is the distribution of (K_b, L_b) conditionally on $\{T_b < +\infty\}$.
If moreover the support of ν is included in $[0; +\infty[$, $w(dk, dl)$ is given explicitly :

$$w(dk, dl) = \frac{-1}{\mathbb{E}(X_1)} \left[\frac{\gamma_0(0)}{2}\, \delta_{0,0}(dk, dl) + (e^{\gamma_0(0)l} - 1) \mathbb{1}_{\{k \geq 0; l \geq 0\}}\, \nu_l(dk)\, dl \right] \tag{4.2}$$

Remark 4.2
1. $\mathcal{N}(0; \sigma^2)$ denotes the Gaussian distribution with mean 0 and variance σ^2.
2. A simular result has been proved by Allan Gut (cf. [7], page 102, Theorem 10.11), where the Lévy process $(X_t\,,\ t \geq 0)$ is replaced by a random walk.

3. a) We observe that time and positions become asymptotically independent. However the two components of the position are not independent. We give a more complete description of the limit distribution of (K_x, L_x) in Proposition 4.4 below.
   b) Obviously $K_x + L_x = X_{T_x} - X_{T_x^-}$ is the jump size of $(X_t\,,\ t \geq 0)$ at T_x. It is easy to deduce from Theorem 4.1 that $\left(\frac{1}{\sqrt{x}}\left(T_x + \frac{x}{\varphi'(-\gamma_0(0))}\right), X_{T_x} - X_{T_x^-}\right)$ converges in distribution to $\mathcal{N}\left(0; -\frac{\varphi''(-\gamma_0(0))}{\varphi'^3(-\gamma_0(0))}\right) \otimes w_0$ where w_0 is the probability measure on $\mathbb{R}_+$:

$$\begin{aligned} w_0(ds) =& \frac{-1}{\mathbb{E}(X_1)}\left[\frac{\gamma_0(0)}{2}\,\delta_0(ds) + \frac{e^{\gamma_0(0)s} - 1 - \gamma_0(0)s}{\gamma_0(0)} 1\!\!1_{\{s \geq 0\}}\,\nu(ds)\right. \\ & \left. + \int_{\mathbb{R}_-}\left(\int_0^{-y}(1 - e^{\gamma_0(0)(b+y)})F(0,0,0,b)n(b,ds)db\right)\nu(dy)1\!\!1_{\{s \geq 0\}}\right], \end{aligned} \tag{4.3}$$

   and $n(b, ds)$ is the distribution of $X_{T_b} - X_{T_b^-}$ conditionally on $\{T_b < +\infty\}$. If the jumps of $(X_t\,,\ t \geq 0)$ are positive, $w_0(ds)$ can be simplified :

$$w_0(ds) = \frac{-1}{\mathbb{E}(X_1)}\left[\frac{\gamma_0(0)}{2}\,\delta_0(ds) + \frac{e^{\gamma_0(0)s} - 1 - \gamma_0(0)s}{\gamma_0(0)} 1\!\!1_{\{s \geq 0\}}\,\nu(ds)\right]. \tag{4.4}$$

4. a) If $\mathbb{E}(X_1) > 0$, then $T_x < +\infty$ a.s. and the triplet $\left(\frac{1}{\sqrt{x}}\left(T_x + \frac{x}{\varphi'(0)}\right), K_x, L_x\right)$ converges in distribution to $\mathcal{N}\left(0; -\frac{\varphi''(0)}{\varphi'^3(0)}\right) \otimes w$, where w is defined by the relation obtained replacing $\gamma_0(0)$ by $-\gamma_0^*(0)$ in (4.1).
   In particular if $(X_t\,,\ t \geq 0)$ has only positive jumps :

$$w(dk, dl) = \frac{1}{\mathbb{E}(X_1)}\left[\frac{\gamma_0^*(0)}{2}\,\delta_{0,0}(dk, dl) + (1 - e^{-\gamma_0^*(0)l})1\!\!1_{\{k \geq 0; l \geq 0\}}\,\nu_l(dk)\,dl\right]. \tag{4.5}$$

   b) In the third case : $\mathbb{E}(X_1) = 0$, the normalization concerning T_x has to be modified. In that case $\left(\frac{T_x}{x^2}, K_x, L_x\right)$ converges in distribution to $\varrho \otimes w$, where ϱ denotes the law of the first hitting time of level $\sqrt{\frac{1}{\varphi''(0)}}$ by a standard Brownian motion started at 0. Recall that $\int_0^{+\infty} e^{-\theta x}\varrho(dx) = e^{-\sqrt{\frac{2\theta}{\varphi''(0)}}}$. Moreover $\varphi''(0) = 1 + \int_{\mathbb{R}} y^2\nu(dy))$.

The probability measure w on $\mathbb{R}_+ \times \mathbb{R}_+$ is defined as follows :

$$\begin{aligned} w(dk, dl) = & \frac{1}{\varphi''(0)} \Big[\delta_{0,0}(dk, dl) + 2l \mathbb{1}_{\{k \geq 0; l \geq 0\}} \, \nu_l(dk) \, dl \\ & -2 \int_{\mathbb{R}_-} \left(\int_0^{-y} (b+y) n(b, dk, dl) db \ \mathbb{1}_{\{k \geq 0, l \geq 0\}} \right) \nu(dy) \Big] \,, \end{aligned} \tag{4.6}$$

where $n(b, dk, dl)$ is the distribution of (K_b, L_b).
This expression may be simplified if $(X_t \, , \, t \geq 0)$ has only positive jumps :

$$w(dk, dl) = \frac{1}{\varphi''(0)} \left[\delta_{0,0}(dk, dl) + 2l \mathbb{1}_{\{k \geq 0; l \geq 0\}} \, \nu_k(dl) \, dk \right] \tag{4.7}$$

Proof of Theorem 4.1
Our approach is based on the following estimate :

$$F(\theta, \mu, \rho, x) \sim_{x \to +\infty} C_0(\theta, \mu, \rho) e^{-\gamma_0(\theta) x} \,, \tag{4.8}$$

where $F(\theta, \mu, \rho, x) = \mathbb{E}\left(e^{-\theta T_x - \mu K_x - \rho L_x} \mathbb{1}_{\{T_x < +\infty\}} \right)$, and $C_0(\theta, \mu, \rho)$ (resp. $\gamma_0(\theta)$) is determined by (2.72), (2.73) (resp. (2.7)).
The three cases $\mathbb{E}(X_1) > 0$, $\mathbb{E}(X_1) < 0$ and $\mathbb{E}(X_1) = 0$ may be treated with the same technic (cf. [12], Introduction, point 7.). Therefore we only deal with the third case : $\mathbb{E}(X_1) = 0$.
For simplicity we restrict ourselves to the couple (T_x, K_x), instead of the triplet (T_x, K_x, L_x).
Since $\varphi(-\gamma_0(\theta)) = \theta$, $\varphi(0) = 0$ and $\varphi'(0) = 0$, taking the asymptotic expansion of φ at 0, of order 2, we obtain :

$$\theta = \varphi(-\gamma_0(\theta)) = \varphi(-h) = \frac{h^2}{2} \varphi''(0) + \mathrm{o}(h^2) \,. \tag{4.9}$$

Consequently :

$$\gamma_0(\theta) = h = \sqrt{\frac{2\theta}{\varphi''(0)}} + \mathrm{o}(\sqrt{\theta}) \,. \tag{4.10}$$

Recall that (4.8) is uniform with respect to $\theta \in [0, \kappa]$. Then choosing $\rho = 0$ and replacing θ by $\dfrac{\theta}{x^2}$ in (4.8) bring to :

$$\mathbb{E}\left(e^{-\theta \frac{T_x}{x^2} - \mu K_x} \right) \sim_{x \to +\infty} C_0(0, \mu) e^{-\sqrt{\frac{2\theta}{\varphi''(0)}}} \,. \tag{4.11}$$

We have to check that $C_0(0, \mu)$ and $e^{-\sqrt{\frac{2\theta}{\varphi''(0)}}}$ are Laplace transforms of probability measures on $[0, +\infty[$. As for $e^{-\sqrt{\frac{1}{\varphi''(0)}}\sqrt{2\theta}}$, it is well known (cf. [8] page 96 formula

(8.6)) that it is the Laplace transform of the first hitting time of level $\sqrt{\frac{1}{\varphi''(0)}}$ by a standard Brownian motion started at 0.
We modify the identity (2.73) (with $\rho = 0$) via $\dfrac{e^{-ay} - 1 + ay}{a^2} = -\displaystyle\int_0^y (z-y)e^{-az}dz$.
Then we obtain :

$$\begin{aligned} C_0(0,\mu) = & \frac{1}{\varphi''(0)} \left[1 + 2\int_0^{+\infty} e^{-\mu z} \left(\int_{[z,+\infty[} (y-z)\nu(dy) \right) dz \right. \\ & \left. -2\int_0^{+\infty} e^{-\mu z} \int_0^{+\infty} db \left(\int_{-\infty}^{b} (y+b)e^{-(y+b)x}\nu(dy) \right) n(b,dz) \right] \\ = & \int_0^{+\infty} e^{-\mu z} w(dz) \,. \end{aligned} \tag{4.12}$$

□

Remark 4.3 *We would like to point out that the asymptotic development of* $F(\theta,\mu,\rho,x)$ *given by (2.23) gives the rate of convergence of* $\left(\frac{1}{\sqrt{x}} \left(T_x + \frac{x}{\varphi'(-\gamma_0(0))} \right) \right)$ *to the Gaussian distribution. Suppose that* $\mathbb{E}[X_1] > 0$*. Let* A_x *be the distribution function of* $\left(\frac{1}{\sqrt{x}} \left(T_x + \frac{x}{\varphi'(-\gamma_0(0))} \right) \right)$ *and* $\widehat{A}_x$ *be its characteristic function :*

$$\begin{aligned} A_x(t) &= \mathbb{P}\left(\frac{1}{\sqrt{x}}\big(T_x + \frac{x}{\varphi'(-\gamma_0(0))}\big) \le t \right), \quad t \in \mathbb{R} \\ \widehat{A}_x(\theta) &= \mathbb{E}\Big[e^{\frac{i\theta}{\sqrt{x}}(T_x + \frac{x}{\varphi'(-\gamma_0(0))})} \Big], \qquad \theta \in \mathbb{R}. \end{aligned}$$

Thanks to (2.23), it is no difficult to check that, if x *is large enough :*

$$\sup_{\theta>0} |\widehat{A}_x(\theta) - \widehat{A}(\theta)| \le \frac{k\theta}{\sqrt{x}}, \tag{4.13}$$

where k *is a positive constant and* $\widehat{A}$ *is the characteristic function of* $\mathcal{N}\left(0; -\frac{\varphi''(-\gamma_0(0))}{\varphi'^3(-\gamma_0(0))}\right)$.
Berry-Essen's inequalities (see for instance [10], p 285) implies the existence of two positives constants k_1 *and* k_2 *such that :*

$$\sup_{t\in\mathbb{R}} |A_x(t) - A(t)| \le k_1 \int_0^a \frac{k\theta}{\sqrt{x}} \frac{d\theta}{\theta} + \frac{k_2}{a}, \quad \text{for any } \ a > 0, \tag{4.14}$$

where A *is the distribution function of the previous Gaussian distribution.*

Choosing $a = x^{1/4}$*, we obtain :*

$$\sup_{t\in\mathbb{R}} |A_x(t) - A(t)| \leq \frac{k_3}{x^{1/4}},\ x \geq 1, \tag{4.15}$$

for some $k_3 > 0$*.*

We would like to provide a stochastic interpretation of the probability measure w defined by (4.2). Let (K, L) be a two-dimensional r.v. with probability distribution w. Obviously $\{K = 0\} = \{L = 0\}$ and this event occurs with probability $\frac{-\gamma_0(0)}{2\mathbb{E}(X_1)}$. Conditionally on $\{L > 0\}$, the distribution of (K, L) is of type $\alpha\left(e^{\gamma_0(0)l} - 1\right)\mathbb{1}_{\{k>0;l>0\}}\nu_l(dk)dl$ where ν_l is the positive measure defined in Theorem 4.1.
This leads us to consider the positive measure :

$$w^*(dk, dl) = \alpha(e^{\gamma l} - 1)\mathbb{1}_{\{k>0;l>0\}}\nu_l(dk)dl\,, \tag{4.16}$$

where $\gamma > 0$, ν_l is the image of ν by $y \to y - l$ and ν is a positive measure on $]0; +\infty[$ satisfying :

$$\int_0^{+\infty} (e^{\gamma k} - 1 - \gamma k)\nu(dk) < +\infty\,, \tag{4.17}$$

α being the normalization factor : $\alpha = \dfrac{\gamma}{\displaystyle\int_0^{+\infty} (e^{\gamma k} - 1 - \gamma k)\nu(dk)}$.

Proposition 4.4 *Let* (K^*, L^*) *be a two dimensional r.v. with distribution* w^* *defined by (4.16) . Then* L^* *has a density function given by* $\alpha(e^{\gamma l} - 1)\nu([l, +\infty[)\mathbb{1}_{\{l>0\}}$*. Conditionally to* $L^* = l$*, the distribution of* $S^* = L^* + K^*$ *is* $\frac{1}{\nu([l, +\infty[)}\mathbb{1}_{\{s>l\}}\nu(ds)$*.*

We have considered w defined by (4.1), but a similar analysis can be developed with w satisfying (4.6).

References

[1] J. Bertoin and R.A. Doney. Cramér's estimate for Lévy processes. *Statistics and Probability Letters*, 21:363–365, 1994.

[2] R.A. Doney. Hitting probabilities for spectrally positive Lévy processes. *J.London Math. Soc.*, 44:566–576, 1991.

[3] F. Dufresne and H.U. Gerber. Risk theory for the compound Poisson process that is perturbed by diffusion. *Insurance: Mathematics and Economics*, 10(1991):51–59, 1990.

[4] A.G. Frolova, Y.M. Kabanov, and S.M. Pergamenshchikov. In the insurance business risky investments are dangerous. *Preprint, Univ. de Franche Comté, Besançon*, 2001.

[5] H.J. Furrer and H. Schmidli. Exponential inequalities for ruin probabilities of risk processes perturbed by diffusion. *Insurance: Mathematics and Economics*, 15:23–36, 1994.

[6] H.U. Gerber and B. Landry. On the discounted penalty at ruin in a jump-diffusion and the perpetual put option. *Insurance: Mathematics and Economics*, 22:263–276, 1998.

[7] A. Gut. *Stopped Random Walks. Limit Theorems and Applications*, volume 5 of *Applied Probability.* Springer-Verlag, New York, 1988.

[8] I. Karatszas and S.E. Shreve. *Brownian motion and Stochastic Calculus.* Springer-Verlag, New York, second edition, 1991. Graduate Texts in Mathematics, 113.

[9] C. Kluppelberg, A.E. Kyprianou, and R.A. Maller. Ruin Probabilities and Overshoot for General Lévy Insurance Risk Processes. preprint, 2003.

[10] M. Loève. *Probability theory.* Third edition. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1963.

[11] J. Paulsen. On Cramèr-like asymptotics for risk processes with stochastic return on investments. *Preprint*, 2001.

[12] B. Roynette, P. Vallois, and A. Volpi. Lévy processes : Hitting time, overshoot and undershoot; I- functional equations. *Preprint*, 2003.

[13] H. Schmidli. Cramèr-Lundberg approximations for ruin probabilities of risk processes perturbed by diffusion. *Insurance: Mathematics and Economics*, 16:135–149, 1995.

[14] H. Schmidli. Distribution of the first ladder height of a stationary risk process perturbed by α-stable Lévy motion. *Insurance: Mathematics and Economics*, 28:13–20, 2001.